

PEPTIDE DERIVATIVES

FIELD OF THE INVENTION

5 The present invention relates to novel derivatives of peptide hormones and analogues thereof which have a protracted profile of action and to methods of making and using them.

10 BACKGROUND OF THE INVENTION

Peptide hormones are widely used in medical practice and since they can be produced by recombinant DNA technology it can be expected that their importance will increase also in the years
15 to come. When native peptide hormones or analogues thereof are used in therapy it is generally found that they have a high clearance rate. A high clearance rate of a therapeutic agent is inconvenient in cases where it is desired to maintain a high blood level thereof over a prolonged period of time since
20 repeated administrations will then be necessary. Examples of peptide hormones which have a high clearance rate are: ACTH, corticotropin-releasing factor, angiotensin, calcitonin, insulin and fragments and analogues thereof, glucagon, glucagon-like peptide and analogues and fragments thereof,
25 IGF-1, IGF-2, enterogastrin, somatostatin, somatotropin, somatomedin, parathyroid hormone, thrombopoietin, erythropoietin, hypothalamic releasing factors, prolactin, thyroid stimulating hormones, endorphins, enkephalins, vasopressin, oxytocin, opioids and analogues thereof,
30 superoxide dismutase, interferon, asparaginase, arginase, arginine deaminase, adenosine deaminase and ribonuclease.

Although it has in some cases been possible to influence the release profile of peptide hormones by applying suitable
35 pharmaceutical compositions this approach has various shortcomings and is not generally applicable. Accordingly, there still is a need for improvements in the field of

administration of peptide hormones.

SUMMARY OF THE INVENTION

5

In the present text, the term peptide is used to designate both small peptides and polypeptides and proteins. The terms peptide and peptide hormone are used as encompassing both naturally occurring and synthetic peptide hormones and
10 fragments and analogues thereof. Analogues are peptides in which one or more amino acids in the parent peptide have been deleted or substituted by another amino acid, or to which one or more amino acids have been added, and which still have qualitatively - but not necessarily quantitatively - the same
15 pharmacological effect as the parent peptide.

The present invention relates generally to novel derivatives of peptide hormones which have a protracted profile of action.

20 Thus, in its broadest aspect, the present invention relates to a pharmacologically active peptide hormone which has been modified by introducing a lipophilic substituent comprising from 8 to 40 carbon atoms in either the N-terminal or the C-terminal amino acid of the native peptide hormone or an
25 analogue thereof, with the proviso that when the lipophilic substituent is attached to the N-terminal amino group then the substituent comprises a group which can be negatively charged and with the further proviso, that said peptide hormone is not insulin or an analogue thereof.

30

In one preferred embodiment of the present invention, a carboxyl group contained in the lipophilic group, W, forms an amide bond together with the α -amino group of the N-terminal amino acid of the parent peptide.

35

In another preferred embodiment of the present invention, a carboxyl group contained in the lipophilic group, W, forms an

amide bond together with the ϵ -amino group of a N-terminal lysine.

In another preferred embodiment of the present invention, the
5 lipophilic group, W, is composed of a spacer and a bulk
lipophilic substituent. The spacer is preferably succinic
acid, Glu or Asp. The bulk lipophilic substituent is
preferably a straight chain fatty acid which optionally has an
10 carboxyl groups forms an amide bond with an amino group in the
N-terminal amino acid of the parent peptide while the other
carboxyl group forms an amide bond with an amino group
contained in the bulk lipophilic group. When Glu or Asp is
used as spacer, one of the carboxyl groups forms an amide bond
15 with an amino group in the N-terminal amino acid of the parent
peptide while the bulk lipophilic substituent preferably is
the acyl group of a straight chain fatty acid or of an acid
which comprises a partly or completely hydrogenated
cyclopentanophenanthrene skeleton which acyl group is attached
20 to the amino group of the spacer.

In another preferred embodiment of the present invention, an
amino group contained in the lipophilic group Z forms an amide
bond together with the carboxyl group of the C-terminal amino
25 acid of the parent peptide.

In another preferred embodiment of the present invention, Z is
a straight chain fatty acid which has an amino group.

30 In another preferred embodiment of the present invention, Z
has a group which can be negatively charged.

In another preferred embodiment of the present invention, Z
has a free carboxylic acid group.

35

In another preferred embodiment of the present invention, the
lipophilic group Z is composed of a spacer and a bulk

lipophilic substituent. The spacer is preferably Lys, Glu or Asp. When Lys is used as spacer, the bulk lipophilic substituent, in one preferred embodiment, is the acyl group of a straight chain fatty acid or of an acid which comprises a partly or completely hydrogenated cyclopentanophenanthrene skeleton which acyl group is attached to the amino group of the spacer group. In a further preferred embodiment, when Lys is used as spacer, a further spacer is inserted between the ϵ -amino group of Lys and the bulk lipophilic substituent. In one preferred embodiment, such a further spacer is succinic acid which forms an amide bond with the ϵ -amino group of Lys and with an amino group present in the bulk lipophilic substituent. In another preferred embodiment such a further spacer is Glu or Asp which form one amide bond with the ϵ -amino group of Lys and a further amide bond with a carboxyl group present in the bulk lipophilic substituent which is preferably a straight chain fatty acid or an acid which comprises a partly or completely hydrogenated cyclopentanophenanthrene skeleton.

In another preferred embodiment, the present invention relates to the use of the peptide derivatives of the invention as medicaments.

In another preferred embodiment, the present invention relates to medicaments containing the peptide derivatives of the invention.

In another preferred embodiment, the present invention relates to a pharmaceutical composition for the treatment of osteoporosis in a patient in need of such a treatment, comprising a therapeutically effective amount of an IGF-1 derivative according to the invention together with a pharmaceutically acceptable carrier.

In another preferred embodiment, the present invention relates to a method of treating osteoporosis in a patient in need of

such a treatment comprising administering to the patient a therapeutically effective amount of an IGF-1 derivative according to the invention together with a pharmaceutically acceptable carrier.

5

Examples of parent peptide hormones which are of interest in connection with the present invention are the following: ACTH, corticotropin-releasing factor, angiotensin, calcitonin, glucagon, glucagon-like peptide and analogues and fragments thereof e.g. GLP-1 and GLP-2 and analogues and fragments thereof, IGF-1, IGF-2, enterogastrin, somatostatin, somatotropin, somatomedin, parathyroid hormone, thrombopoietin, erythropoietin, hypothalamic releasing factors, prolactin, thyroid stimulating hormones, endorphins, enkephalins, vasopressin, oxytocin, opioids and analogues thereof, superoxide dismutase, interferon, asparaginase, arginase, arginine deaminase, adenosine deaminase and ribonuclease.

20 Examples of particularly preferred derivatives of IGF-1 and IGF-1 analogues are:

Lys⁶⁸ (N^ε-tetradecanoyl) des(69,70) human IGF-1;

Lys⁶⁸ [N^ε-γ-Glu(N^α-hexadecanoyl)-OH]-OH des(69,70) human IGF-1;

25 Lys⁶⁹ (N^ε-tetradecanoyl) des(70) human IGF-1;

Ser⁶⁹-NH(CH₂)_nCOOH des(70) human IGF-1 wherein n is an integer from 12 to 24;

Ser⁶⁹-NH(CH₂)_nCH₃ des(70) human IGF-1 wherein n is an integer from 12 to 24;

30 Lys⁷¹ (N^ε-tetradecanoyl) human IGF-1;

Ala⁷⁰-NH(CH₂)_nCOOH human IGF-1 wherein n is an integer from 12 to 24; and

Ala⁷⁰-NH(CH₂)_nCH₃ human IGF-1 wherein n is an integer from 12 to 24.

35

A preferred derivative of somatostatin is:

Ala-Gly-Cys-Lys-Asn-Phe-Phe-Trp-Lys-Thr-Tyr-Thr-Ser-Cys-Lys [N^{ϵ} - γ -Glu(N^{α} -tetradecanoyl)-OH]-OH (the two Cys residues are connected via a disulphide bridge).

- 5 A preferred derivative of GLP-1 is:

His-Ala-Glu-Gly-Thr-Phe-Thr-Ser-Asp-Val-Ser-Ser-Tyr-Leu-Glu-Gly-Gln-Ala-Ala-Lys-Glu-Phe-Ile-Ala-Trp-Leu-Val-Lys [N^{ϵ} - γ -Glu(N^{α} -tetradecanoyl)-OH]-OH.

10

A preferred ANP analogue is:

Ser-Leu-Arg-Arg-Ser-Ser-Cys-Phe-Gly-Gly-Arg-Met-Asp-Arg-Ile-Gly-Ala-Gln-Ser-Gly-Leu-Gly-Cys-Asn-Ser-Phe-Arg-Tyr-Lys [N^{ϵ} - γ -Glu(N^{α} -tetradecanoyl)-OH]-OH.

15

A preferred type of derivative of a dynorphin analogue is:

Tyr-Gly-Gly-Phe-Cys-Arg-Arg-D-Ala-Arg-Pro-Cys-NH-(CH₂)_n-COOH, wherein n is an integer from 8 to 24.

- 20 A preferred derivative of enterogastrin is:

H-Ala-Pro-Gly-Pro-Arg-Lys (N^{ϵ} -tetradecanoyl)-OH.

DETAILED DESCRIPTION OF THE INVENTION

25

Pharmaceutical compositions

- Pharmaceutical compositions containing a peptide derivative according to the present invention may be administered parenterally to patients in need of such a treatment. Parenteral administration may be performed by subcutaneous, intramuscular or intravenous injection by means of a syringe, optionally a pen-like syringe. Alternatively, parenteral administration can be performed by means of an infusion pump.
- 30 A further option is a composition which may be a powder or a liquid for the administration of the peptide derivative in the

form of a nasal spray. As a still further option, it may also be possible to administer the peptide derivatives transdermally.

5 Pharmaceutical compositions containing a compound of the present invention may be prepared by conventional techniques, e.g. as described in Remington's Pharmaceutical Sciences, 1985.

10 Thus, the injectable compositions of the peptide derivatives of the invention can be prepared using the conventional techniques of the pharmaceutical industry which involves dissolving and mixing the ingredients as appropriate to give the desired end product.

15 Thus, according to one procedure, the peptide derivative is dissolved in an amount of water which is somewhat less than the final volume of the composition to be prepared. An isotonic agent, a preservative and a buffer is added as
20 required and the pH value of the solution is adjusted - if necessary - using an acid, e.g. hydrochloric acid, or a base, e.g. aqueous sodium hydroxide as needed. Finally, the volume of the solution is adjusted with water to give the desired concentration of the ingredients.

25 Examples of isotonic agents are sodium chloride, mannitol and glycerol.

30 Examples of preservatives are phenol, m-cresol, methyl p-hydroxybenzoate and benzyl alcohol.

Examples of suitable buffers are sodium acetate and sodium phosphate.

35 A composition for nasal administration of certain peptide hormones may, for example, be prepared as described in European Patent No. 272097 (to Novo Nordisk A/S).

The peptide derivatives of this invention can be used in the treatment of various diseases. The particular peptide derivative to be used and the optimal dose level for any patient will depend on the disease to be treated and on a variety of factors including the efficacy of the specific peptide derivative employed, the age, body weight, physical activity, and diet of the patient, on a possible combination with other drugs, and on the severity of the case. It is recommended that the dosage of the peptide derivative of this invention be determined for each individual patient by those skilled in the art in a similar way as for known peptide hormones.

The present invention is further illustrated by the following examples which, however, are not to be construed as limiting the scope of protection. The features disclosed in the foregoing description and in the following examples may, both separately and in any combination thereof, be material for realizing the invention in diverse forms thereof.

EXAMPLES

25 Abbreviations:

| | | | |
|----|------|---|--|
| | Fmoc | : | 9-fluorenylmethyloxycarbonyl. |
| | For | : | formyl |
| | Dde | : | 1-(4,4-dimethyl-2,6-dioxocyclohexylidene)-ethyl. |
| 30 | DMF | : | N,N-dimethylformamide. |
| | Tbu | : | <i>tert</i> -butyl. |
| | Acm | : | acetamidomethyl. |
| | DIC | : | N,N'-diisopropylcarbodiimide. |
| | HOBT | : | 1-hydroxybenzotriazole. |
| 35 | TFA | : | trifluoroacetic acid. |

Analytical

Molecular masses of the products prepared were obtained by plasma desorption mass spectrometry (PDMS) using Bio-Ion 20 instrument (Bio-Ion Nordic AB, Uppsala, Sweden).

Determination of lipophilicity.

The lipophilicity of peptides and peptide derivatives relative to human insulin, k'_{rel} , was measured on a LiChrosorb RP18 (5 μ m, 4x250 mm) HPLC column by isocratic elution at 40°C using mixtures of A) 0.1 M sodium phosphate buffer, pH 7.3, containing 10% acetonitrile, and B) 50% acetonitrile in water as eluents. The elution was monitored by following the UV absorption of the eluate at 214 nm. Void time, t_0 , was found by injecting 0.1 mM sodium nitrate. Retention time for human insulin, $t_{insulin}$, was adjusted to at least $2t_0$ by varying the ratio between the A and B solutions. $k'_{rel} = (t_{derivative} - t_0) / (t_{insulin} - t_0)$.

EXAMPLE 1

Synthesis of For-Nle-Leu-Phe-Nle-Tyr-Lys(N^ε-tetradecanoyl)-OH.

25

For-Nle-Leu-Phe-Nle-Tyr-Lys-OH, was purchased from Bachem Feinchemikalien AG, Switzerland. The peptide is a potent chemoattractant for human neutrophils. The title compound was prepared by dissolving 17 mg of For-Nle-Leu-Phe-Nle-Tyr-Lys-OH in 5 ml of DMF and then adding 35 μ l of triethylamine followed by 20 mg of solid tetradecanoic acid succinimidyl-N-hydroxy ester to the solution. The reaction was monitored by RP-HPLC employing a column packed with reversed phase C18 silica material. For the elution was used a gradient from 30% ethanol to 80 % ethanol in 0.1% aqueous TFA. The product was purified on a column (length 250 mm diameter 20 mm) packed with C18

silica reversed phase material. The compound was dissolved in 74% ethanol/0.1% aqueous TFA and subsequently applied to the column and purified at 40 °C by isocratic elution in the same buffer at a flow rate of 6 ml/hour. The yield was 20 mg. The identity of the compound was confirmed by PDMS.

Molecular mass, found by PDMS: 1034, theory: 1034.

The lipophilicity of the title compound relative to human insulin was found to be 8.2×10^3 .

Reference

The reference compound, For-Nle-Leu-Phe-Nle-Tyr-Lys-OH, was purchased from Bachem Feinchemikalien AG, Switzerland, and used as received. The lipophilicity of the reference compound relative to human insulin was found to be 2.3.

EXAMPLE 2

Synthesis of H-Tyr-D-Ala-Gly-Phe-Leu-Lys(N^εtetradecanoyl)-OH.

The enkephalin derivative H-Tyr-D-Ala-Gly-Phe-Leu-Lys(N^ε-tetradecanoyl)-OH was made from Boc-Tyr-D-Ala-Gly-Phe-Leu-Lys-OH (A-2435 Bachem Feinchemikalien AG, Switzerland). The Boc-Tyr-D-Ala-Gly-Phe-Leu-Lys-OH was acylated using tetradecanoic acid succinimidyl-N-hydroxy ester as described in Example 1. The reaction mixture was evaporated to dryness and the residue was dissolved in TFA and evaporated to dryness, solubilized in ethanol/water/0.1% and purified by RP-HPLC as described in Example 1. The yield was 15 mg.

Molecular mass, found by PDMS: 909, theory: 907.

The lipophilicity of the title compound relative to human insulin was found to be 2.3×10^3 .

Reference

The reference compound, H-Tyr-D-Ala-Gly-Phe-Leu-Lys-OH, was synthesized from Boc-Tyr-D-Ala-Gly-Phe-Leu-Lys-OH by dissolving 20 mg of this compound in 200 μ l of TFA and evaporating to dryness. The residue was dissolved in 5% acetic acid and freeze dried. The lipophilicity of the reference compound relative to human insulin was found to be 3.0×10^{-3} .

10 **EXAMPLE 3**

Synthesis of H-Pro-His-Pro-Phe-His-Phe-Phe-Val-Tyr-Lys(N^{ϵ} -tetradecanoyl)-OH.

15 Fmoc-Pro-His-Pro-Phe-His-Phe-Phe-Val-Tyr-Lys-OH (obtained from Bachem Feinchemikalien AG, Switzerland) which is a potent inhibitor of renin was allowed to react with tetradecanoic acid succinimidyl-N-hydroxy ester as described in Example 1. After the acylation reaction, the Fmoc group was removed by
20 addition of piperidine to the reaction mixture to a final concentration of 20%. The title compound was isolated by RP-HPLC as described in Example 1. The yield was 23 mg.

Molecular mass, found by PDMS: 1529.6, theory: 1529.

25

The lipophilicity relative to human insulin was found to be 5.3×10^{-3} .

Reference

30 The reference compound, H-Pro-His-Pro-Phe-His-Phe-Phe-Val-Tyr-Lys-OH, was synthesized from Fmoc-Pro-His-Pro-Phe-His-Phe-Phe-Val-Tyr-Lys-OH (obtained from Bachem Feinchemikalien AG, Switzerland). Thus, 20 mg of Fmoc-Pro-His-Pro-Phe-His-Phe-Phe-Val-Tyr-Lys-OH was dissolved in 500 μ l of 20% piperidine in
35 DMF and left for 20 min. The reference compound was purified by RP-HPLC as described in Example 1.

The lipophilicity of the reference compound relative to human insulin was found to be 2.3×10^2 .

5 **EXAMPLE 4**

Synthesis of Arg⁴,Arg⁹,Lys¹⁵(N⁸-tetradecanoyl) somatostatin.

10 The title compound was synthesized from Fmoc-Arg⁴,Arg⁹,Lys¹⁵ somatostatin which was obtained from Saxon Biochemicals GMBH, Hannover, Germany. 50 mg of Fmoc-Arg⁴,Arg⁹,Lys¹⁵ somatostatin was dissolved in a mixture of 346 μ l of DMF and 53.9 μ l of 4-methylmorpholine. The mixture was cooled to 15 °C and 15.9 mg of tetradecanoic acid succinimidyl-N-hydroxy ester dissolved
15 in 100 μ l of DMF was added. The reaction was allowed to proceed for 3 hours and 20 min and then stopped by addition of 4140 μ l of 5% acetic acid in DMF. The title compound was purified by RP-HPLC as follows: The sample was applied to a column (10x250 mm) of Lichrosorb RP-18 (7 μ m) Merck, Germany,
20 Art. 9394. The column was equilibrated with a mixture of 90% buffer A (50 mM tris, 75 mM (NH₄)₂SO₄ adjusted to pH 7.0 with H₂SO₄, 20% CH₃CN) and 10% of buffer B (80% CH₃CN). The sample was applied to the column and eluted with a linear gradient from 10% to 90 % of buffer B in buffer A at a flow rate of 4
25 ml/hour at 40 °C. The fractions containing the title compound were evaporated to dryness, dissolved in 50% acetic acid and desalted by gel filtration at 4 °C employing of column (16x150 mm) of BIO GEL P2 (BIO RAD, California, USA). The fractions containing the desired product were diluted with water and
30 freeze dried. The yield was 2 mg. The identity of the compound was confirmed by PDMS.

Molecular mass, found by PDMS: 2033, theory 2032.

35

Determination of protraction in pigs

The title peptide derivative of Example 4 was ^{125}I -labelled with Boulton & Hunters reagent (Bolton, A.E. and Hunter, W.M. (1973) Biochem. J. **133**. 529-539) as follows : 50 nmol of peptide was dissolved in 1 ml of DMSO and subsequently 400 μl of DMF and 2 μl of N-ethylisopropylamine were added. The solution was added to an amount of Boulton & Hunters reagent containing 500 μCi of radioactivity. The reaction was allowed to proceed for 20 min. and then 10 μl of ethanolamine in DMF was added. The polypeptide was purified and isolated by RP-HPLC employing a column (4x250 mm) at a flow rate of 1 ml/min as described above.

As a measure of the protraction, the disappearance rate in pigs was studied and $T_{50\%}$ was determined. $T_{50\%}$ is the time when 50% of the ^{125}I -labelled peptide has disappeared from the site of injection as measured with an external γ -counter (Ribet, U et al., The Pig as a Model for Subcutaneous Absorption in Man. In: M. Serrano-Rios and P.J. Lefebvre (Eds): Diabetes 1985; Proceedings of the 12th Congress of the International Diabetes Federation, Madrid, Spain, 1985 (Excerpta Medica, Amsterdam, (1986) 891-96).

Subcutaneous injection of the ^{125}I -labelled peptide derivative in pigs showed a $T_{50\%}$ of 1.7 ± 0.5 h (n=4), whereas the non tetradecanoylated, ^{125}I -labelled reference peptide showed a $T_{50\%}$ of 0.7 ± 0.1 h.

Reference

The ^{125}I -labelled reference peptide was synthesized from Fmoc-Arg⁴,Arg⁹,Lys¹⁵ somatostatin. Thus, 20 mg of Fmoc-Arg⁴,Arg⁹,Lys¹⁵ somatostatin was dissolved in 1000 μl of 20% piperidine/DMF. After 20 min the product was purified by RP-HPLC, desalted and freeze dried and labelled with Boulton & Hunters reagent as described in Example 4.

EXAMPLE 5

Synthesis of Lys¹⁵(N⁶-tetradecanoyl) atrial natriuretic peptide.

5 Human (H-Ser-Leu-Arg-Arg-Ser-Ser-Cys-Phe-Gly-Gly-Arg-Met-Asp-Arg-Ile-Gly-Ala-Gln-Ser-Gly-Leu-Gly-Cys-Asn-Ser-Phe-Arg-Tyr-Lys(N⁶-tetradecanoyl)-COOH) was synthesized by standard Fmoc solid phase peptide synthesis (Methods in Molecular Biology, 10 Vol 35: Peptide Synthesis Protocols). The ε-amino group of the C-terminal lysine was acylated using tetradecanoic acid succinimidyl-N-hydroxy ester according to the procedure described below. The synthesis was performed manually in polypropylene syringes, on a resin based on a low cross linked polystyrene 15 backbone grafted with polyoxyethylene (TentaGel Resin).

Procedure:

One gram of resin was added 3 equivalents of the acid labile 20 linker 4-hydroxymethylphenoxyacetic acid (HMPA). 3 equivalents of Fmoc-Lys(Dde)-OH was coupled as the first amino acid, with 0.5 equivalent of 4-dimethylaminopyridine as activating reagent. The Fmoc-protecting group was cleaved with 20% piperidine/DMF for 30 minutes. All other amino acids were 25 coupled as N^α-Fmoc protected amino acids with a mixture of DIC/HOBT (1:1 eq) in DMF as activating reagents. The amino acid Cys, was coupled as Fmoc-Cys(Acm)-OH. The cysteines were deprotected and oxidized by treatment with 10 Mm Iodine in DMF for 2 minutes. After the last Fmoc-protecting group was 30 removed, the N^α-group of the last coupled amino acid was protected with the Boc group by coupling with 5 equivalents of di-*tert*-butyl-dicarbonate. The Dde-protecting group of N⁶-Lys was cleaved with 2% hydrazine/DMF for 20 minutes, and the free N⁶-group was acylated with 5 equivalents of tetradecanoic acid 35 succinimidyl-N-hydroxy ester. The Boc-, *t*Bu-protecting groups and the HMPA-linker were cleaved with 95% TFA/5% H₂O for 1.5

hour. The TFA/H₂O was evaporated under reduced pressure, and the peptide was precipitated in diethyl ether as the HCl-salt, and freeze dried from a 10 mM ammonium hydrogen carbonate (pH 8.8). The overall yield was 35 mg. By N-terminal sequencing
5 the product was shown to have the correct sequence.

Molecular mass, found by PDMS: 3417, which corresponds to the calculated mass plus sodium.

10

EXAMPLE 6

Lys³⁰(N^ε-decanoyl) glucagon.

15

The title compound was purchased from Saxon Biochemicals GMBH, Hannover, Germany, as custom synthesis.

4.32 mg Lys³⁰(N^ε-decanoyl), glucagon (equivalent to 4 mg glucagon) was dissolved in 4 ml of 1.8 mM hydrochloric acid
20 added 0.9% sodium chloride and pH of the solution was measured to 2.7. The resulting solution was sterilized by filtration and transferred to a vial.

Two groups of rabbits (n=6 in each) were injected with 2
25 IU/animal of Insulin Actrapid at time -60 min. At time t=0 group 1 was injected SC with molar equivalent of 0.54 mg of Lys³⁰(N^ε-decanoyl) glucagon/rabbit and group 2 injected SC with 0.5 mg of glucagon/rabbit. Blood was sampled at times: -60, 0, 15, 30, 60, 120, 180 and 240 min, and the glucose concentration determined by the hexokinase method. The resulting
30 blood glucose concentrations are given in the table in mg glucose/100 ml:

| min | -60 | 0 | 15 | 30 | 60 | 120 | 180 | 240 |
|----------|-----|----|----|-----|-----|-----|-----|-----|
| glucagon | 98 | 49 | 93 | 102 | 111 | 94 | 88 | 67 |

| | | | | | | | | |
|------------------------|----|----|----|----|-----|-----|-----|-----|
| glucagon derivative | 94 | 51 | 79 | 93 | 114 | 112 | 116 | 110 |
|------------------------|----|----|----|----|-----|-----|-----|-----|

As it appears from the table, the blood glucose raising effect of glucagon is retained in Lys³⁰(N₈-decanoyl) glucagon but with a prolonged action.